



Les effets du dérangement liés aux activités humaines sur la faune sauvage

Synthèse bibliographique

Travail réalisé par Aurélie Bisch, stagiaire au Parc naturel régional des Ballons des Vosges en 2018.

- Arlettaz, R., Patthey, P., Baltic, M., Leu, T., Schaub, M., Palme, R., & Jenni-Eiermann, S. (2007). Spreading free-riding snow sports represent a novel serious threat for wildlife. *Proceedings of the Royal Society B: Biological Sciences*, 274(1614), 1219–1224. <https://doi.org/10.1098/rspb.2006.0434>
- Baines, D., & Richardson, M. (2007). An experimental assessment of the potential effects of human disturbance on Black Grouse Tetrao tetrix in the North Pennines , England, 149, 56–64.
- Baltic, M., Jenni-Eiermann, S., Arlettaz, R., & Palme, R. (2005). A noninvasive technique to evaluate human-generated stress in the black grouse. *Annals of the New York Academy of Sciences*, 1046, 81–95. <https://doi.org/10.1196/annals.1343.008>
- Braunisch, V., & Suchant, R. (2007). A Model for Evaluating the “Habitat Potential” of a Landscape for Capercaillie Tetrao urogallus: A Tool for Conservation Planning. *Wildlife Biology*, 13(sp1), 21–33. [https://doi.org/10.2981/0909-6396\(2007\)13\[21:AMFETH\]2.0.CO;2](https://doi.org/10.2981/0909-6396(2007)13[21:AMFETH]2.0.CO;2)
- Braunisch, V., & Suchant, R. (2013). Aktionsplan Auerhuhn Tetrao urogallus im Schwarzwald : Ein integratives Konzept zum Erhalt einer überlebensfähigen Population, 41, 29–41.
- BRENOT, J. ., CATUSSE, M., & MENONI, E. (1996). Effets de la station de ski de fond du Plateau de Beille (Ariège) sur une importante population de Grand Tétras.
- Graf, R. ., Bollmann, K., Suter, W., & Bugmann, H. (2004). Using a multi-scale model for identifying priority areas in capercaillie (, 84–90.
- Green, R., & Higginbottom, K. (2000). The effects of non-consumptive wildlife tourism on free-ranging wildlife: a review. *Pacific Conservation Biology*, 6(3), 183–197. <https://doi.org/10.1071/PC000183>
- Knight, R. L., & Gutzwiler, K. J. (1995). *Wildlife and recreationists: Coexistence through management and research*. *Wildlife and Recreationists: Coexistence Through Management and Research*. Retrieved from

<http://books.google.com/books?hl=en&lr=&id=BRbBAvLwQIAC&oi=fnd&pg=PR2&dq=Wildlife+and+Recreationists+Coexistence+through+management+and+research&ots=tLxWRTHxBC&sig=tGeL7wgT3G4a6dZOTZo3gFr7Lg>

Lanc, R. B., Uillemain, M. G., Ouronval, J. M., Esmonts, D. D., & Ritz, H. F. (2006). Effects of non-consumptive leisure disturbance to wildlife, 61, 117–134. Retrieved from http://documents.irevues.inist.fr/bitstream/handle/2042/55674/RevuedEcologie_2006_61_2_117.pdf?sequence=1&isAllowed=y

Marshall, K. (2005). Capercaillie and recreational disturbance study, (January), 50.

Ménoni, E., Favre-Ayala, V., Cantegrel, R., Revenga, J., Camprodón, J., García, D., Campion, D. et Riba, L. (2012). Réflexion technique pour la prise en com pte du Grand tétras dans la gestion forestière pyrénéenne.

Mollet, P., & Thiel, D. (2009). Wintertourismus beeinflusst das Verhalten und die Stressphysiologie des Auerhuhns | Winter tourism influences the behaviour and the stress physiology of the capercaillie. *Schweizerische Zeitschrift Fur Forstwesen*, 160(10), 311–317. <https://doi.org/10.3188/szf.2009.0311>

Montadert, M. (2013). Tétras lyre et dérangement touristique : Synthèse bibliographique, 49.

Moss, R., & Picozzi, N. (1994). Management of Forests for Capercaillie in Scotland, 33. Retrieved from [http://www.forestry.gov.uk/pdf/FCBU113.pdf/\\$FILE/FCBU113.pdf](http://www.forestry.gov.uk/pdf/FCBU113.pdf/$FILE/FCBU113.pdf)

Perrin, N., & Neet, C. (2003). Winter habitat selection by two sympatric forest grouse in western Switzerland : implications for conservation, 112, 373–382.

Sachot, S., & Neet, C. (2006). GESTION SYLVICOLE ET GRAND TÉTRAS : LES ACTIONS DU CANTON DE VAUD (SUISSE), 62, 60–62. <https://doi.org/10.1051/nss:2006057>

Sachot, S., Perrin, N., & Neet, C. (2006). Viability and management of an endangered Capercaillie (*Tetrao urogallus*) metapopulation in the Jura Mountains, western Switzerland. *Biodiversity and Conservation*, 15(6), 2017–2032. <https://doi.org/10.1007/s10531-005-0771-y>

SANIGA, M. (2003). Causes of the population decline in capercaillie (*Tetrao urogallus*) in the West Carpathians.

Segelbacher, G., Höglund, J., & Storch, I. (2003). From connectivity to isolation: Genetic consequences of population fragmentation in capercaillie across Europe. *Molecular Ecology*, 12(7), 1773–1780. <https://doi.org/10.1046/j.1365-294X.2003.01873.x>

Stettler, M., & Christen, R. (2010). Vielfältiger Lebensraum, Ruhe, Sensibilisierung – Auerhuhnförderung auf drei Ebenen. *Schweizerische Zeitschrift Für Forstwesen*, 161(7), 258–263. <https://doi.org/10.3188/szf.2010.0258>

Storch, I. (2013). Human disturbance of grouse - why and when? *Wildlife Biology*, 19(4), 390–403. <https://doi.org/10.2981/13-006>

Stratégie nationale d'actions en faveur du grand tétras *Tetrao urogallus major*. (2012).

Suárez-Seoane, S., & García-Rovés, P. (2004). Do disturbances in surrounding areas affect a core population of Cantabrian capercaillie *Tetrao urogallus cantabricus*? The case of the The Natural Reserve of Muniellos (Asturias, NW Spain). *Ardeola*, 51(2), 395–409.

Suchant, R., & Schäfer, A. (2002). Integrating Tourism and Grouse Habitat Protection in the Black Forest, 95–101.

Summers, R. W., Proctor, R., Thornton, M., & Avey, G. (2004). Habitat selection and diet of the Capercaillie *Tetrao urogallus* in Abernethy Forest, Strathspey, Scotland. *Bird Study*,

51(1), 58–68. <https://doi.org/10.1080/00063650409461333>

Thiel, D., Jenni-Eiermann, S., Braunisch, V., Palme, R., & Jenni, L. (2008). Ski tourism affects habitat use and evokes a physiological stress response in capercaillie *Tetrao urogallus*: A new methodological approach. *Journal of Applied Ecology*, 45(3), 845–853. <https://doi.org/10.1111/j.1365-2664.2008.01465.x>

Thiel, D., Jenni-Eiermann, S., Palme, R., & Jenni, L. (2011). Winter tourism increases stress hormone levels in endangered capercaillie *Tetrao urogallus*. *Ibis*, 153, 122–133.

Thiel, D. K. (2007). Behavioral and physiological effects in Capercaillie (*Tetrao urogallus*) caused by human disturbance. *Journal of Wildlife Management*. Retrieved from <https://www.bioone.org/doi/full/10.2193/2006-268>

THIEL, D., MÉNONI, E., BRENOT, J.-F., & JENNI, L. (2007). Effects of Recreation and Hunting on Flushing Distance of Capercaillie. *Journal of Wildlife Management*, 71(6), 1784–1792. <https://doi.org/10.2193/2006-268>

Vogelwarte, & University of Bern. (2008). Coqs de bruyère : prière de ne pas déranger ! *Agriculture*.